



Handbook for

- GEL SV
- PCR SV
- CLEANUP SV
- COMBO GP

expin™

Expin™ Protocol Handbook  
For purification of fragment DNA

# Customer & Technical Support

Do not hesitate to ask us any question.

We thank you for any comment or advice.

## Contact us at

[www.geneall.com](http://www.geneall.com)

Tel : 82-2-407-0096

Fax : 82-2-407-0779

E-mail(Order/Sales) : [sales@geneall.com](mailto:sales@geneall.com)

E-mail(Tech. Info.) : [tech@geneall.com](mailto:tech@geneall.com)

## Visit GeneAll® Community

[www.geneall.com](http://www.geneall.com)

[www.geneall.co.kr](http://www.geneall.co.kr)

## Trademarks

AmpONE™, DirEx™, Exfection™, Exgene™, Expin™, Exprep™, EzClear™, EzSep™, GenEx™, Hybrid-Q™, RiboEx™, Riboclear™, Ribospin™ are trademarks of GeneAll Biotechnology co., ltd.

© 2013 GeneAll® Biotechnology, all right reserved.

This protocol handbook is included in :

GeneAll® Expin™ Gel SV (102-150, 102-102)

GeneAll® Expin™ PCR SV (103-150, 103-102)

GeneAll® Expin™ CleanUp SV (113-150, 113-102)

GeneAll® Expin™ Combo GP (112-150, 112-102)

Visit [www.geneall.com](http://www.geneall.com) or [www.geneall.co.kr](http://www.geneall.co.kr) for FAQ, QnA and more information.



# INDEX

## GeneAll® Expin™ Protocols

Kit Contents	4
Precautions and Disclaimer	5
Storage Conditions	5
Quality Control	5
Safety Information	5
Introduction	6
Product Specifications	7
Applicable range in length of DNA	7
General considerations	8
Binding	8
Washing	9
Elution	9
Correlation between the elution volume and the recovery	10
Electrophoresis analysis	11
<b>PROTOCOLS FOR..</b>	
Gel SV Protocol	12
PCR SV Protocol	18
CleanUp SV Protocol	22
Troubleshooting Guide	26
Ordering Information	28
Expin™ Gel SV Brief protocol	30
Expin™ PCR SV Brief protocol	30
Expin™ CleanUp SV Brief protocol	31



## KIT CONTENTS

Cat. No.	Gel SV		PCR SV	
	102-150	102-102	103-150	103-102
<b>Size</b>	<b>mini</b>	<b>mini</b>	<b>mini</b>	<b>mini</b>
No. of preparation	50	200	50	200
GeneAll® SV Column type D	50	200	50	200
Collection Tube	50	200	50	200
Buffer GB	60 ml	120 ml x 2	-	-
Buffer PB	-	-	30 ml	120 ml
Buffer NW	60 ml	250 ml	60 ml	250 ml
Buffer EB*	15 ml	30 ml	15 ml	30 ml
Protocol Handbook	1	1	1	1

Cat. No.	CleanUp SV		Combo GP	
	113-150	113-102	112-150	112-102
<b>Size</b>	<b>mini</b>	<b>mini</b>	<b>mini</b>	<b>mini</b>
No. of preparation	50	200	50	200
GeneAll® SV Column type D	50	200	50	200
Collection Tube	50	200	50	200
Buffer GB	-	-	60 ml	120 ml x 2
Buffer PB	-	-	30 ml	120 ml
Buffer NR	30 ml	120 ml	-	-
Buffer NW	60 ml	250 ml	60 ml	250 ml
Buffer EB*	15 ml	30 ml	15 ml	30 ml
Protocol Handbook	1	1	1	1

\* 10 mM TrisCl, pH 8.5



## PRECAUTIONS AND DISCLAIMER

GeneAll® Expin™ Kit is for research use only, and should not be used for drug, household or other unintended uses. All due care and attention should be taken in every procedure in this handbook. Please consult Material Safety Data Sheet (MSDS) for information regarding hazards and safe handling practices.



## STORAGE CONDITIONS

All components of GeneAll® Expin™ Gel, PCR and CleanUp SV Kit should be stored at room temperature (15~25°C) and are stable for one year. It should be protected from exposure to direct sunlight.

During shipment or storage under cold ambient condition, a precipitate may be formed in buffer GB, PB or NR. In such a case, heat the bottle at 50°C to dissolve completely. Using precipitated buffers will lead to poor DNA recovery. GeneAll® Expin™ SV kit series are guaranteed until the expiration date printed on the product label.



## QUALITY CONTROL

All components in GeneAll® Expin™ Gel, PCR and CleanUp SV Kit are manufactured in strictly clean condition, and its degree of cleanness is monitored periodically. Restriction enzyme assay, PCR amplification assay, spectrophotometric assay and automatic sequencing as the validation of quality are carried out from lot to lot thoroughly, and only the qualified is approved to be delivered.



## SAFETY INFORMATION

The buffers included in GeneAll® Expin™ Gel, PCR and CleanUp SV Kit contain irritant which is harmful when in contact with skin or eyes, or when inhaled or swallowed. Care should be taken during handling. Always wear gloves and eye protector, and follow standard safety precautions.

Buffer GB, PB and NR contain chaotropes. It can form highly reactive compounds when combined with bleach. Do NOT add bleach or acidic solutions directly to the sample-preparation waste.

# Introduction

GeneAll® Expin™ SV series provide reliable and fast methods for the purification of fragment DNA from agarose gel and PCR/enzyme reaction mixtures. GeneAll® Expin™ SV series consist of;

## **GeneAll® Expin™ Gel SV (102-150, 102-102)**

Gel SV kit is designed for a fast and efficient isolation of 80 bp to 10 kb of DNA fragments from standard or low-melting agarose gel in TAE or TBE buffer system. Purified DNA can be directly used in ligation, labelling, sequencing and many other downstream application without further manipulation.

## **GeneAll® Expin™ PCR SV (103-150, 103-102)**

PCR SV kit provides a simple and rapid method to purify PCR products or other enzymatic reactions in just 6 minutes. Up to 10 ug of pure DNA (100 bp ~ 10 kb) can be obtained, and this purified DNA can be directly used in cloning, sequencing and many other application. PCR SV procedures remove the DNA fragment smaller than 100 bp, resulting in removal of primers and primer dimers in PCR products.

## **GeneAll® Expin™ CleanUp SV (113-150, 113-102)**

CleanUp SV kit provides a simple and rapid method to purify 40 bp ~ 10 kb DNA from enzymatic reactions in just 6 minutes. Up to 10 ug of pure DNA which is at least 40 bp but less than 10 kb in length can be obtained using this kit and the purified DNA can be directly used for sequencing, cloning and other routine applications without further manipulation.


## **GeneAll® Expin™ Combo GP (112-150, 112-102)**

The combination set of GeneAll® Gel SV and PCR SV.

# Product Specifications

GeneAll® Expin™	Gel SV	PCR SV	CleanUp SV
Standard sample size	200 mg of gel (400 mg max.)	100 ul of PCR reactions	50 ul of enzyme mixtures
Recovered DNA size	80 bp ~ 10 kb	100 bp ~ 10 kb	40 bp ~ 10 kb
Typical yields	~ 85 %	~ 95 %	~ 95 %
Maximum binding capacity	10 ug	10 ug	10 ug
Preparation time	~ 15 mins	~ 6 mins	~ 6 mins
Minimum elution volume	30 ul	30 ul	30 ul

## Applicable range in length of DNA



GeneAll® Expin™ SV series consists of Gel, PCR and CleanUp SV Kit. Each kit is optimized for efficient recovery of DNA and removal of contaminants in each specific application.

## General considerations

GeneAll® Expin™ SV series takes advantage of silica membrane and spin/vacuum column technology to recover DNA fragments. Under high salt conditions DNA binds to silica membrane and impurities pass through membrane into a collection tube. The membranes are washed with an ethanol-containing buffer to remove any traces of proteins, salts, remnant of agarose and other enzymatic reaction components. Finally pure DNA is released into a clean collection tube with water or low ionic strength buffer.

## Binding

The basic principle which DNA binds to silica membrane is identical in all GeneAll® Expin™ SV series. Binding buffers (Buffer GB, PB and NR) make the optimal binding condition in each specific applications. Buffer GB in Gel SV kit is composed to dissolve standard agarose gel as well as low melt agarose gel in addition to adjusting a binding condition. Usually low melt agarose gel results in a better recovery yield. DNA binds to silica membrane at lower pH than pH 7.5. The components and pH of starting sample can alter the pH of the mixture with binding buffer, especially in extraction of DNA from agarose gel. Buffer GB contains pH indicator in order to check this alteration of binding condition. If the color (yellow) of binding mixture turns to brown or purple after addition of buffer GB, it means that the pH of binding mixtures is higher than the optimal, and it can be easily adjusted with small volume of sodium acetate before proceeding with the protocol. The indicator dye is completely removed during subsequent washing steps and does not interfere the downstream applications.



## Washing





Any unwanted oligos and impurities, such as salts, proteins, nucleotides, agarose, dyes and detergents will not be bound but be passed through the silica membrane. A minute impurities, such as salts, are quantitatively washed away with buffer NW which contains ethanol. The quality of DNA can be slightly increased with the repeat of washing. Any residual ethanol should be removed completely with an additional centrifugation because the residual ethanol in eluate may interfere some subsequent applications.

## Elution

DNA is released under the condition of low salts and neutral or weakly alkaline pH ( $7 < \text{pH} < 9.5$ ). Although buffer EB (10 mM TrisCl, pH 8.5), TE, or distilled water can be used for elution, it should be considered that EDTA in buffer TE may interfere the subsequent reactions and low pH ( $< 7.0$ ) of distilled water can reduce DNA recovery. Because water does not have any buffering agents the eluate in water should be stored under  $-20^{\circ}\text{C}$  not to degrade.

The minimum elution volume is 30  $\mu\text{l}$  and lower volume will decrease the yield significantly. It is important for optimal elution to apply the elution buffer to the center of the membrane, because the membrane should be covered completely by eluent for an optimal recovery. Up to 200  $\mu\text{l}$  of elution buffer can be applied and it results in low concentration of DNA. Higher concentrated DNA will be obtained with lower elution volume, and maximum yield can be obtained by larger elution volume. The yield with large fragments ( $> 5$  kb) can be increased slightly by using pre-warmed ( $70^{\circ}\text{C}$ ) elution buffer. Incubation for 1 minute after addition of eluent may increase the efficiency of elution.

# Correlation between the elution volume and the recovery




Correlation between the elution volume and the recover rates. Upper panel; GeneAll® PCR SV Kit. Lower panel; GeneAll® Gel SV Kit. 3 ug of 494 bp PCR products were purified and eluted with the indicated volumes of buffer EB. Elution volume lower than 30 ul causes significant loss of DNA. 1/10 volume of eluate purified with Gel SV Kit was resolved on 1 % agarose gel; lower right. (1/20 for PCR SV Kit, upper right)


DNA size (bp)	Recovery rates (%)		
	Gel SV	PCR SV	CleanUp SV
60	39	0	63
120	71	78	80
200	76	83	84
800	84	94	94
1800	82	91	93
4300	78	85	88
8700	73	76	79

Average recovery rates of GeneAll® Expin™ SV Kit with various sizes of DNA. 3 ug of starting sample was purified and eluted with 50 ul of buffer EB. Optional steps were not performed and SV columns were incubated for 1 minutes after addition of buffer EB.

## Electrophoresis analysis



Quantities of extracted 4.5 kb DNA fragment corresponding to 1/5 of the DNA obtained by purification from 0.5 ug starting DNA with a recovery of 90 %. Sample were run on 1 % TAE agarose gel. M; Lambda-BstP1 marker Lane1; Total amount before extraction (0.5 ug) Lane2; 1/5 amount after extraction [90 ng compared to known amount (10 ~ 150 ng) DNA] \* Total obtained amount of DNA =  $90 \times 5 = 450$  ng approximately (90 %)



PCR products of several sizes (70, 176, 757 and 1487 bp from left to right) were purified using GeneAll® CleanUp SV Kit. The band intensity of 1/5 amount of 20 ul PCR products (lane 1, 5, 9, 13) was almost identical with that of 1/5 amount of 50 ul eluate. (2 ~ 4, 6 ~ 8, 10 ~ 12 and 14 ~ 16, triplicate) M; 1 kb ladder marker

# Expin™ Gel SV Protocol

## Before experiment

- \* All centrifugation should be carried out at 10,000 xg above (> 12,000 rpm) at room temperature in a microcentrifuge.
- \* Prepare water bath or heating block to 50°C.
- \* All solutions should be equilibrated at room temperature before procedures.
- \* For large fragments (>5 kb), pre-warm buffer EB to 70°C.
- \* If Buffer GB has precipitated, dissolve it completely at 50°C before starting.

## Spin/Vacuum Methods for Gel Extraction

When using this kit, one of the two methods can be chosen to purify DNA. DNA fragment can be purified using centrifugation to pull the mixture through the SV column. Alternatively, vacuum can be used to force the mixture out through the column.

## A. Centrifugation Protocol

### 1. Excise the DNA band of interest using a ethanol-cleaned razor blade or scalpel on a transilluminator.

Minimize gel volume by cutting the gel slice as small as possible.

Use of long wave length transilluminator and short handling time will lead to better quality of DNA. It can be critical in some experiments, such as ligation.

### 2. Weigh the gel slice in a microcentrifuge tube. Add 3 volumes (ul) of Buffer GB to 1 volume (mg) of gel.

For example, add 300 ul of buffer GB to each 100 mg of agarose gel slice. For > 1.5 % agarose gel, add 5 volumes of buffer GB.

**3. Incubate at 50°C until the agarose gel is completely melted (5 ~ 10 min).**

To help the efficient dissolving of gel, vortex the tube every 2 ~ 3 min during the incubation.

**4. After the slice has dissolved completely, check that the color of the mixture is yellow (similar to Buffer GB).**

If the color of the mixture becomes brown or purple, add 10 ul of 3M sodium acetate, pH 5.0, and mix. The color of the mixture will turn to yellow. Adjusting of pH is not needed when the change of color is due to the ingredient of loading dye (eg. bromophenol blue, xylene cyanol).

**5. (Optional :) Add 1 gel volume of isopropanol to the sample and vortex to mix.**

For 100 mg of gel volume, add 100 ul of isopropanol. Do NOT centrifuge at this step.

This step is required to increase the recover yields of DNA fragments <200 bp or >5 kb. For the DNA fragments between 200 bp and 5 kb, it has little effect on the recovery.

**6. Transfer the mixture to a SV column. Centrifuge for 1 min. Discard the pass-through and re-insert the SV column into the Collection Tube.**

If the mixture volume is larger than 700 ul, apply the mixture twice; apply 700 ul of the mixture, spin down, discard the pass-through, re-insert empty collection tube, and repeat the step again until all of the mixture has been applied to the SV column.

**7. (Optional :) Apply 500 ul of Buffer GB to the SV column. Centrifuge for 30 sec. Discard the pass-through and reinsert the SV column into the Collection Tube.**

This step is for further complete removal of any traces of agarose and required only for direct use of purified DNA for very sensitive applications, such as in vitro transcription. Usually this step can be skipped for automatic sequencing or ligation.

**8. Add 700 ul of Buffer NW to the SV column. Centrifuge for 30 sec. Discard the pass-through. Reinsert the SV column into the Collection Tube.**

If the purified DNA will be used for salt sensitive applications, let the SV column stand for 5 min after addition of buffer NW, making some amount of wash buffer flow through the column by gravity before centrifugation.

**9. Centrifuge for an additional 1 min to remove residual wash buffer. Transfer the SV column to a new 1.5 ml tube.**

If residual ethanol remains in the SV column, centrifuge again for an additional 1 min at full speed before transferring to a new 1.5 ml tube. Residual ethanol from buffer NW can inhibit subsequent enzymatic reaction.

**10. Apply 50 ul of Buffer EB or ddH<sub>2</sub>O to the center of the membrane in the SV column, let stand for 1 min, and centrifuge for 1 min.**

Ensure that the buffer EB or distilled water is dispensed directly onto the center of SV column membrane for optimal elution of DNA.

To obtain more concentrated DNA solution, apply 30 ul of elution buffer, but the volume lower than 30 ul will decrease the yield significantly. Up to 200 ul of elution buffer can be applied to SV column, and it will reduce the concentration of DNA.

For larger fragment (>5 kb), use pre-warmed (70°C) elution buffer for best efficiency.

For long-term storage, eluting in buffer EB (10 mM TrisCl, pH 8.5) or TE, pH 8.0 and storing under -20°C is recommended. When using water for elution, check the pH of water (>pH 7.0) before elution.

## B. Vacuum Protocol

The vacuum pressure should be in the range of the list below. Lower vacuum pressure may reduce DNA yield and purity.

Most commercial vacuum manifold with luer connectors can be used with this protocol.

15 ~ 18 in Hg	380 ~ 460 mbar
285 ~ 345 mm Hg	5.5 ~ 6.5 Psi

### 1. Excise the DNA band of interest using a ethanol-cleaned razor blade or scalpel on a transilluminator.

Minimize gel volume by cutting the gel slice as small as possible.

Use of long wave length transilluminator and short handling time will lead to better quality of DNA. It can be critical in some experiments, such as ligation.

### 2. Weigh the gel slice in a microcentrifuge tube. Add 3 volumes (ul) of Buffer GB to 1 volume (mg) of gel.

For example, add 300 ul of buffer GB to each 100 mg of agarose gel slice. For > 1.5 % agarose gel, add 5 volumes of buffer GB.

### 3. Incubate at 50°C until the agarose gel is completely melted (5 ~ 10 min).

To help the efficient dissolving of gel, vortex the tube every 2 ~ 3 min during the incubation.

### 4. After the slice has dissolved completely, check that the color of the mixture is yellow (similar to Buffer GB).

If the color of the mixture becomes brown or purple, add 10 ul of 3M sodium acetate, pH 5.0, and mix. The color of the mixture will turn to yellow. Adjusting of pH is not needed when the change of color is due to the ingredient of loading dye (eg. bromophenol blue, xylene cyanol)

**5. (Optional :) Add 1 gel volume of isopropanol to the sample and vortex to mix.**

For 100 mg of gel volume, add 100 ul of isopropanol. Do NOT centrifuge at this step.

This step is required to increase the recover yields of DNA fragments <200 bp or >5 kb. For the DNA fragments between 200 bp and 5 kb, it has little effect on the recovery.

**6. Attach the SV column to a port of the vacuum manifold tightly.**

Most commercial vacuum manifold with luer connectors can be adopted to this protocol.

**7. Transfer the mixture to a SV column, by pipetting. Switch on vacuum source to draw the solution through the SV column. When all liquid has been pulled through the SV column, release the vacuum. Repeat this step until all mixture have applied to the SV column.**

**8. (Optional :) Apply 500 ul of Buffer GB to the SV column and switch on vacuum source. When all liquid has been pulled through the SV column, release the vacuum.**

This step is for further complete removal of any traces of agarose and required only for direct use of purified DNA for very sensitive applications, such as in vitro transcription. Usually this step can be skipped for automatic sequencing.

**9. Apply 700 ul of Buffer NW and switch on vacuum source. When all liquid has been pulled through the SV column, release the vacuum. Transfer the SV column to a empty collection tube (provided).**

If the purified DNA will be used for salt sensitive applications, let the SV column stand for 5 minutes after addition of buffer NW before applying vacuum.

**10. Centrifuge for an additional 1 min to remove residual wash buffer. Transfer the SV column to a new 1.5 ml tube.**

If the SV column has buffer NW associated with it, centrifuge again for additional 1 min at full speed before transferring to a new 1.5 ml tube. Residual ethanol from buffer NW can inhibit subsequent enzymatic reaction.



**11. Apply 50 ul of Buffer EB or ddH<sub>2</sub>O to the center of the membrane in the SV column, let stand for 1 min, and centrifuge for 1 min.**

Ensure that the buffer EB or distilled water is dispensed directly onto the center of SV column membrane for optimal elution of DNA.

To obtain more concentrated DNA solution, apply 30 ul of elution buffer, but the volume lower than 30 ul will decrease the yield significantly. Up to 200 ul of elution buffer can be applied to SV column, and it will reduce the concentration of DNA.

For larger fragment (>5 kb), use pre-warmed (70°C) elution buffer for best efficiency.

For long-term storage, eluting in buffer EB (10 mM TrisCl, pH 8.5) or TE, pH 8.0 and storing under -20°C is recommended. When using water for elution, check the pH of water (>pH 7.0) before elution.

# Expin™ PCR SV Protocol

## Before experiment

- \* All centrifugation should be carried out at 10,000 xg above (> 12,000 rpm) at room temperature in a microcentrifuge.
- \* All solutions should be equilibrated at room temperature before procedures.
- \* For large fragments (>5 kb), pre-warm buffer EB to 70°C.

## Spin/Vacuum Methods for PCR Purification

When using this kit, one of the two methods can be chosen to purify DNA. DNA fragment can be purified using centrifugation to pull the mixture through the SV column. Alternatively, vacuum can be used to force the mixture through the column.

## A. Centrifugation Protocol

- 1. Add 5 volumes of Buffer PB to 1 volume of the sample and mix. Transfer the mixture to a SV column.**  
For 100 ul reaction, add 500 ul of buffer PB. It is not necessary to remove mineral oil.
- 2. Centrifuge for 30 sec. Discard the pass-through and reinsert the SV column back into the same tube.**
- 3. Apply 700 ul of Buffer NW. Centrifuge for 30 sec. Discard the pass-through and reinsert the SV column back into the collection tube.**

**4. Centrifuge for an additional 1 min to remove residual wash buffer. Transfer the SV column to a new 1.5 ml tube.**

If the SV column has buffer NW associated with it, centrifuge again for additional 1 min at full speed before transferring to the new 1.5 ml tube. Residual ethanol from buffer NW can inhibit subsequent enzymatic reaction.

**5. Apply 50 ul of Buffer EB or ddH<sub>2</sub>O to the center of the membrane in the SV column, let stand for 1 min, and centrifuge for 1 min.**

Ensure that the buffer EB or distilled water is dispensed directly onto the center of SV column membrane for optimal elution of DNA.

To obtain more concentrated DNA solution, apply 30 ul of elution buffer, but the volume lower than 30 ul will decrease the yield significantly. Up to 200 ul of elution buffer can be applied to SV column, and it will reduce the concentration of DNA.

For larger fragment (>5 kb), use pre-warmed (70°C) elution buffer for best efficiency.

For long-term storage, eluting in buffer EB (10 mM TrisCl, pH 8.5) or TE, pH 8.0 and storing under -20°C is recommended. When using water for elution, check the pH of water (>pH 7.0) before elution.

## B. Vacuum Protocol

The vacuum pressure should be in the range of the list below. Lower vacuum pressure may reduce DNA yield and purity.

Most commercial vacuum manifold with luer connectors can be used with this protocol.

15 ~ 18 in Hg	380 ~ 460 mbar
285 ~ 345 mm Hg	5.5 ~ 6.5 Psi

- 1. Attach the SV column to a port of the vacuum manifold tightly.**
- 2. Add 5 volumes of Buffer PB to 1 volume of the sample and mix. Transfer the mixture to a SV column by pipetting.**  
For 100 ul reaction, add 500 ul of buffer PB.  
It is not necessary to remove mineral oil.
- 3. Switch on vacuum source to draw the solution through the SV column. When all liquid has been pulled through the SV column, release the vacuum.**
- 4. Apply 700 ul of Buffer NW and switch on vacuum source. When all liquid has been pulled through the SV column, release the vacuum. Transfer the SV column to a collection tube (provided).**
- 5. Centrifuge for 1 min to remove residual wash buffer. Transfer the SV column to a new 1.5 ml tube.**

If the SV column has buffer NW associated with it, centrifuge again for additional 1 min at full speed before transferring to the new 1.5 ml tube. Residual ethanol from buffer NW can inhibit subsequent enzymatic reaction.

**6. Apply 50 ul of Buffer EB or ddH<sub>2</sub>O to the center of the membrane in the SV column, let stand for 1 min, and centrifuge for 1 min.**

Ensure that the buffer EB or distilled water is dispensed directly onto the center of SV column membrane for optimal elution of DNA.

To obtain more concentrated DNA solution, apply 30 ul of elution buffer, but the volume lower than 30 ul will decrease the yield significantly. Up to 200 ul of elution buffer can be applied to SV column, and it will reduce the concentration of DNA.

For larger fragment (>5 kb), use pre-warmed (70°C) elution buffer for best efficiency.

For long-term storage, eluting in buffer EB (10 mM TrisCl, pH 8.5) or TE, pH 8.0 and storing under -20°C is recommended. When using water for elution, check the pH of water (>pH 7.0) before elution.

# Expin™ CleanUp SV Protocol

## Before experiment

- \* All centrifugation should be carried out at 10,000 xg above (> 12,000 rpm) at room temperature in a microcentrifuge.
- \* All solutions should be equilibrated at room temperature before procedures.
- \* For large fragments (>5 kb), pre-warm buffer EB to 70°C.

## Spin/Vacuum Methods for DNA Clean-up

When using this kit, one of the two methods can be chosen to purify DNA. DNA fragment can be purified using centrifugation to pull the mixture through the SV column. Alternatively, vacuum can be used to force the mixture through the column.

## A. Centrifugation Protocol

- 1. Add 10 volumes of Buffer NR to 1 volume of the sample and mix. Transfer the mixture to a SV column.**  
For 50 ul reaction, add 500 ul of buffer NR.  
If the length of DNA is longer than 100 bp, add 5 volumes of buffer NR.
- 2. Centrifuge for 30 sec. Discard the pass-through and reinsert the SV column back into the collection tube.**
- 3. Apply 700 ul of Buffer NW. Centrifuge for 30 sec. Discard the pass-through and reinsert the SV column back into the collection tube.**

**4. Centrifuge for an additional 1 min to remove residual wash buffer. Transfer the SV Column to a new 1.5 ml tube.**

If the SV column has buffer NW associated with it, centrifuge again for additional 1 min at full speed before transferring to the new 1.5 ml tube. Residual ethanol from buffer NW can inhibit subsequent enzymatic reaction.

**5. Apply 50 ul of Buffer EB or ddH<sub>2</sub>O to the center of the membrane in the SV column, let stand for 1 min, and centrifuge for 1 min.**

Ensure that the buffer EB or distilled water is dispensed directly onto the center of SV column membrane for optimal elution of DNA.

To obtain more concentrated DNA solution, apply 30 ul of elution buffer, but the volume lower than 30 ul will decrease the yield significantly. Up to 200 ul of elution buffer can be applied to SV column, and it will reduce the concentration of DNA.

For larger fragment (>5 kb), use pre-warmed (70°C) elution buffer for best efficiency.

For long-term storage, eluting in buffer EB (10 mM TrisCl, pH 8.5) or TE, pH 8.0 and storing under -20°C is recommended. When using water for elution, check the pH of water (>pH 7.0) before elution.

## B. Vacuum Protocol

The vacuum pressure should be in the range of the list below. Lower vacuum pressure may reduce DNA yield and purity.

Most commercial vacuum manifold with luer connectors can be used with this protocol.

15 ~ 18 in Hg	380 ~ 460 mbar
285 ~ 345 mm Hg	5.5 ~ 6.5 Psi

- 1. Attach the SV column to a port of the vacuum manifold tightly.**
- 2. Add 10 volumes of Buffer NR to 1 volume of the sample and mix. Transfer the mixture to a SV column by pipetting.**  
For 50 ul reaction, add 500 ul of buffer NR.  
If the length of DNA is longer than 100 bp, add 5 volumes of buffer NR.
- 3. Switch on vacuum source to draw the solution through the SV column. When all liquid has been pulled through the SV column, release the vacuum.**
- 4. Apply 700 ul of Buffer NW and switch on vacuum source. When all liquid has been pulled through the spin column, release the vacuum. Transfer the SV column to a collection tube (provided).**
- 5. Centrifuge for an additional 1 min to remove residual wash buffer. Transfer the SV column to a new 1.5 ml tube.**  
If the SV column has buffer NW associated with it, centrifuge again for additional 1 min at full speed before transferring to the new 1.5 ml tube. Residual ethanol from buffer NW can inhibit subsequent enzymatic reaction.



**6. Apply 50 ul of Buffer EB or ddH<sub>2</sub>O to the center of the membrane in the SV column, let stand for 1 min, and centrifuge for 1 min.**

Ensure that the buffer EB or distilled water is dispensed directly onto the center of SV column membrane for optimal elution of DNA.

To obtain more concentrated DNA solution, apply 30 ul of elution buffer, but the volume lower than 30 ul will decrease the yield significantly. Up to 200 ul of elution buffer can be applied to SV column, and it will reduce the concentration of DNA.

For larger fragment (>5 kb), use pre-warmed (70°C) elution buffer for best efficiency.

For long-term storage, eluting in buffer EB (10 mM TrisCl, pH 8.5) or TE, pH 8.0 and storing under -20°C is recommended. When using water for elution, check the pH of water (>pH 7.0) before elution.

## Troubleshooting Guide

Facts	Possible Causes	Suggestions
<b>Low or no recovery</b>	<b>[Gel SV] Incompletely solubilized gel</b>	<p>The sliced agarose gel should be completely dissolved without any particles. To assist the complete solubilization, mix the tube by vortexing every 2 ~ 3 minutes during incubation, or increase the incubation time. To use low melt agarose usually results better recovery.</p>
	<b>[Gel SV] Too high pH of binding mixture</b>	<p>At high pH, the binding of DNA to silica membrane will be significantly reduced. The dye included in buffer GB indicates the pH of mixture as color change from yellow at optimal pH to brown or purple at abnormally higher pH. If the color of mixture has turned to brown or purple, add 10 ul of 3M sodium acetate, pH 5.0 to the sample and mix. The color of mixture will turn to yellow indicating the correct pH for DNA binding.</p>
	<b>Improper elution buffer</b>	<p>As user's requirement, elution buffer other than buffer EB can be used. The condition of optimal elution is low salt concentration with alkaline pH (<math>7 &lt; \text{pH} &lt; 9</math>). When water or other buffer was used as eluent, ensure that conditions.</p>
	<b>Elution buffer incorrectly dispensed</b>	<p>Ensure that elution buffer dispensed to the center of membrane. Incorrectly dispensed elution buffer causes inappropriate contact with membrane, followed by poor DNA recovery.</p>
<b>Ligation failure</b>	<b>[Gel SV] Too long or strong exposure to UV on transilluminator</b>	<p>UV destroys the DNA ends. Use UV of long wave length and make the handling time as short as possible when excising the gel slice.</p>

Facts	Possible Causes	Suggestions
<b>Column clogging</b>	<b>[Gel SV] Incompletely solubilized gel</b>	See the section 'Incompletely solubilized gel' in the Facts "Low or no recovery"
	<b>[Gel SV] &gt; 1.5% agarose gel is used</b>	<p>For &gt; 1.5 % agarose gel, 5 volumes of buffer GB to 1 volume of gel slice should be added. For 100 mg of agarose gel, add 500 ul of buffer GB.</p> <p>If the column is clogged, transfer the mixture from the SV column to a 1.5 ml tube, add 1 volume of buffer GB to mixture volume. Incubate for 5 minutes at 50°C, proceed again to binding steps.</p>
<b>Enzymatic reaction is not performed well with the purified DNA</b>	<b>Residual ethanol from Buffer NW remains in eluate</b>	It is essential to remove any residual ethanol included in buffer NW from column membrane. Centrifuge again for complete removal of ethanol.
	<b>Too high salt concentration in eluate</b>	Incubate for 5 minutes after addition of buffer NW at washing steps.
	<b>Eluate contains denatured ssDNA</b>	For reannealing of ssDNA to dsDNA, incubate ssDNA at 95°C for 2 min, and then allow to cool slowly to room temperature.
<b>DNA floats out of well while loading of agarose gel</b>	<b>Residual ethanol from Buffer NW remains in eluate</b>	It is essential to remove any residual ethanol included in buffer NW from SV column membrane. Centrifuge again for complete removal of ethanol.
<b>Nonspecific band appears after purification</b>	<b>DNA denatured</b>	Renature the DNA by warming up to 95°C for a minute and let cool slowly to room temperature.

# Ordering Information

Products	Scale	Size	Cat. No.	Type
----------	-------	------	----------	------

## GeneAll® Hybrid-Q™ for rapid preparation of plasmid DNA

Plasmid Rapidprep	50	100-150	mini / spin
	100	100-102	

## GeneAll® Exprep™ for preparation of plasmid DNA

Plasmid SV	mini	50	101-150	spin / vacuum
		200	101-102	
		1,000	101-111	
	Midi	26	101-226	spin / vacuum
		50	101-250	
		100	101-201	

## GeneAll® Exfection™ for preparation of highly pure plasmid DNA

Plasmid LE (Low Endotoxin)	mini	50	111-150	spin / vacuum	
		200	111-102		
	Midi	26	111-226	spin / vacuum	
		100	111-201		
	Plasmid EF (Endotoxin Free)	Midi	20	121-220	spin
			100	121-201	

## GeneAll® Expin™ for purification of fragment DNA

Gel SV	mini	50	102-150	spin / vacuum
		200	102-102	
PCR SV	mini	50	103-150	spin / vacuum
		200	103-102	
CleanUp SV	mini	50	113-150	spin / vacuum
		200	113-102	
Combo GP	mini	50	112-150	spin / vacuum
		200	112-102	

## GeneAll® Exgene™ for isolation of total DNA

Tissue SV	mini	100	104-101	spin / vacuum
		250	104-152	
	Midi	26	104-226	spin / vacuum
		100	104-201	
	MAXI	10	104-310	spin / vacuum
		26	104-326	
Tissue plus! SV	mini	100	109-101	spin / vacuum
		250	109-152	
	Midi	26	109-226	spin / vacuum
		100	109-201	
	MAXI	10	109-310	spin / vacuum
		26	109-326	

Products	Scale	Size	Cat. No.	Type
----------	-------	------	----------	------

## GeneAll® Exgene™ for isolation of total DNA

Blood SV	mini	100	105-101	spin / vacuum	
		250	105-152		
	Midi	26	105-226	spin / vacuum	
		100	105-201		
	MAXI	10	105-310	spin / vacuum	
		26	105-326		
Cell SV	mini	100	106-101	spin / vacuum	
		250	106-152		
	MAXI	10	106-310	spin / vacuum	
		26	106-326		
	Clinic SV	mini	100	108-101	spin / vacuum
			250	108-152	
Midi		26	108-226	spin / vacuum	
		100	108-201		
MAXI		10	108-310	spin / vacuum	
		26	108-326		
Genomic DNA micro	mini	50	118-050	spin / vacuum	
		100	117-101		
	Midi	26	117-226	spin / vacuum	
		100	117-201		
	MAXI	10	117-310	spin / vacuum	
		26	117-326		
Soil DNA mini	mini	50	114-150	spin	
		100	114-102		
GMO SV	mini	50	107-150	spin / vacuum	
		200	107-102		

## GeneAll® GenEx™ for isolation of total DNA

GenEx™ Blood	Sx	100	220-101	solution
		500	220-105	
	Lx	100	220-301	solution
		500	221-101	
GenEx™ Cell	Sx	100	221-101	solution
		500	221-105	
	Lx	100	221-301	solution
		500	222-101	
GenEx™ Tissue	Sx	100	222-101	solution
		500	222-105	
	Lx	100	222-301	solution

Products	Scale	Size	Cat. No.	Type
----------	-------	------	----------	------

**GeneAll® GenEx™** for isolation of total DNA

GenEx™ Plant	Sx	100	227-101	solution
	Mx	100	227-201	
	Lx	100	227-301	
GenEx™ Plant plus!	Sx	100	228-101	solution
	Mx	50	228-250	
	Lx	20	228-320	

**GeneAll® DirEx™**  
for preparation of PCR-template without extraction

DirEx™		50	250-050	solution
--------	--	----	---------	----------

**GeneAll® RNA series** for preparation of total RNA

RiboEx™	mini	100	301-001	solution
		200	301-002	
Hybrid-R™	mini	100	305-101	spin
Hybrid-R™ Blood RNA	mini	50	315-150	spin
Hybrid-R™ miRNA	mini	50	325-150	spin
RiboEx™ LS	mini	100	302-001	solution
		200	302-002	
Riboclear™	mini	50	303-150	spin
Riboclear™ plus!	mini	50	313-150	spin
Ribospin™	mini	50	304-150	spin
Ribospin™ vRD	mini	50	302-150	spin
Ribospin™ vRD plus!	mini	50	312-150	spin
Ribospin™ Plant	mini	50	307-150	spin
Allspin™	mini	50	306-150	spin

**GeneAll® AmpONE™** for PCR amplification

Taq DNA polymerase	250 U	501-025	(2.5 U/μℓ)
	500 U	501-050	
	1,000 U	501-100	
α-Taq DNA polymerase	250 U	502-025	(2.5 U/μℓ)
	500 U	502-050	
	1,000 U	502-100	
Pfu DNA polymerase	250 U	503-025	(2.5 U/μℓ)
	500 U	503-050	
	1,000 U	503-100	

Products	Scale	Size	Cat. No.	Type
----------	-------	------	----------	------

**GeneAll® AmpONE™** for PCR amplification

Hotstart Taq DNA polymerase	250 U	531-025	(2.5 U/μℓ)	
	500 U	531-050		
	1,000 U	531-100		
Clean Taq DNA polymerase	250 U	551-025	(2.5 U/μℓ)	
	500 U	551-050		
	1,000 U	551-100		
Clean α-Taq DNA polymerase	250 U	552-025	(2.5 U/μℓ)	
	500 U	552-050		
	1,000 U	552-100		
Taq Premix	20 μℓ	521-200	lyophilized	
	50 μℓ	521-500		
	20 μℓ	526-200	solution	
50 μℓ	526-500			
α-Taq Premix	20 μℓ	522-200	lyophilized	
	50 μℓ	522-500		
	20 μℓ	527-200	solution	
50 μℓ	527-500			
HS-Taq Premix	20 μℓ	525-200	solution	
	50 μℓ	525-500		
Taq Premix (w/o dye)	96 tubes	20 μℓ	524-200	lyophilized
α-Taq Premix (w/o dye)	96 tubes	20 μℓ	525-200	solution
dNTP mix		500 μℓ	509-020	2.5 mM each
dNTP set (set of dATP, dCTP, dGTP and dTTP)		1 ml x 4 tubes	509-040	100 mM

**GeneAll® AmpMaster™** for PCR amplification

Taq Master mix	2x	541-010	0.5 ml x 2 tubes
	2x	541-050	0.5 ml x 10 tubes
α-Taq Master mix	2x	542-010	0.5 ml x 2 tubes
	2x	542-050	0.5 ml x 10 tubes
HS-Taq Master mix	2x	545-010	0.5 ml x 2 tubes
	2x	545-050	0.5 ml x 10 tubes

\* Each dNTP is available

## GeneAll® Expin™ Gel SV Brief protocol

*\* Before use this protocol, we strongly recommend you read first carefully the detailed protocol at page 12.*

Prepare a water bath or a heating block to 50°C.

Check if precipitates appear in buffer GB. If so, dissolve completely at 50°C before proceeding.

1. Excise the DNA band of interest as small and rapid as possible.
2. Add 3 vol (ul) of buffer GB to 1 vol (mg) of gel.
3. Incubate at 50°C with occasional mixing until the gel is completely melted.  
⇒ Check that the color of mixture is yellow after melting. If not, see the step 4 in detailed protocol at page 13.
4. Transfer the mixture to a SV column, centrifuge for 1 min, discard the filtrate.
5. Add 700 ul of buffer NW to the SV column, centrifuge for 30 sec and discard the filtrate.
6. Centrifuge for an additional 1 min and transfer the SV column to a new 1.5 ml tube.
7. Apply 50 ul of buffer EB or ddH<sub>2</sub>O, let stand for 1 min and centrifuge for 1 min.

## GeneAll® Expin™ PCR SV Brief protocol

*\* Before use this protocol, we strongly recommend you read first carefully the detailed protocol at page 18.*

1. Add 5 vol of buffer PB to 1 vol of the sample and mix thoroughly.
2. Transfer the mixture to a SV column, centrifuge for 30 sec and discard the filtrate.
3. Add 700 ul of buffer NW to the SV column, centrifuge for 30 sec and discard the filtrate.
4. Centrifuge for an additional 1 min and transfer the SV column to a new 1.5 ml tube.
5. Apply 50 ul of buffer EB or ddH<sub>2</sub>O, let stand for 1 min and centrifuge for 1 min.

# GeneAll® Expin™ CleanUp SV Brief protocol

*\* Before use this protocol, we strongly recommend you read first carefully the detailed protocol at page 22.*

Check if precipitates appear in buffer NR. If so, dissolve completely at 50°C before proceeding.

1. Add 10 (5) vol of buffer NR to 1 vol of the sample and mix thoroughly.  
⇒ If the length of target DNA is larger than 100 bp, use 5 vol of buffer NR.
2. Transfer the mixture to a SV column, centrifuge for 30 sec and discard the filtrate.
3. Add 700 ul of buffer NW to the SV column, centrifuge for 30 sec and discard the filtrate.
4. Centrifuge for an additional 1 min and transfer the SV column to a new 1.5 ml tube.
5. Apply 50 ul of buffer EB or ddH<sub>2</sub>O, let stand for 1 min and centrifuge for 1 min.



**[www.geneall.com](http://www.geneall.com)**

GeneAll Bldg., 303-7 Dongnam-ro  
Songpa-gu, Seoul, Korea 138-859

E-mail : [sales@geneall.com](mailto:sales@geneall.com)

Tel : 82-2-407-0096

Fax : 82-2-407-0779

©2013 GeneAll Biotechnology, All right reserved

Edited by **BnP**  
Designed by **Park Eun Ah**